


Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims

1. (Currently Amended) A compound of formula (I):



(I)

wherein:

Ring A is pyridin-2-yl or thiazol-2-yl; wherein said pyridin-2-yl or thiazol-2-yl ~~is~~ may be optionally substituted on carbon by one or more groups selected from R^4 ;

~~One~~ one of R^1 and R^2 is hydrogen and the other is hydrogen or C_{1-4} alkyl; wherein R^1 and R^2 are optionally ~~may be~~ substituted on carbon by one or more groups selected from R^5 ;

R^3 is selected from C_{1-4} alkyl, C_{1-4} alkoxy, carbocyclyl, heterocyclyl, carbocycloxy and heterocycloxy; wherein R^3 ~~is~~ may be independently optionally substituted on carbon by one or more groups selected from R^6 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen ~~may be~~ is optionally substituted by C_{1-4} alkyl;

R^4 is selected from halo, carboxy and C_{1-4} alkyl;


R^5 and R^6 are independently selected from halo, C_{1-4} alkyl, C_{1-4} alkoxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, carbocyclyl, heterocyclyl, carbocycloxy, heterocycloxy and carbocyclylidenyl; wherein R^5 and R^6 ~~may be~~ are independently optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen ~~may be~~ is optionally substituted by C_{1-4} alkyl;

R^7 is selected from halo, carboxy, methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, dimethylamino, diethylamino and N -methyl- N -ethylamino;

or a salt, solvate or pro-drug thereof.

2. (Original) A compound according to Claim 1 wherein Ring A is unsubstituted or is substituted by carboxy.
3. (Currently Amended) A ~~compounds~~ compound according to ~~any one of the preceding claims~~ Claim 1 wherein one of R^1 and R^2 is hydrogen and the other is hydrogen or C_{1-4} alkyl.
4. (Currently Amended) A compound according to ~~any one of the preceding claims~~ Claim 1 wherein R^3 is selected from C_{1-4} alkoxy; wherein R^3 ~~may be~~ is independently optionally substituted on carbon by one or more groups selected from R^6 .
5. (Currently Amended) A compound according to ~~any one of the preceding claims~~ Claim 1 wherein R^3 is selected from 2-fluorobenzyloxy, 5-methylisoxazol-3-ylmethoxy and 2-thien-3-ylethoxy
6. (Original) A compound according to Claim 1 selected from:
 - 2-methyl-4-isobutoxy-6-[*N*-(5-carboxypyridin-2-yl)carbamoyl]benzofuran;
 - 2-methyl-4-(2-fluorophenylmethoxy)-6-[*N*-(5-carboxypyridin-2-yl)carbamoyl]benzofuran;
 - 2-methyl-4-isobutoxy-6-[*N*-(5-carboxythiazol-2-yl)carbamoyl]benzofuran;
 - 2-methyl-4-(5-methylisoxazol-3-ylmethoxy)-6-[*N*-(5-carboxypyridin-2-yl)carbamoyl]benzofuran;
 - 4-(2-fluorophenylmethoxy)-6-[*N*-(5-carboxypyridin-2-yl)carbamoyl]benzofuran;
 - 4-(5-methylisoxazol-3-ylmethoxy)-6-[*N*-(5-carboxypyridin-2-yl)carbamoyl]benzofuran;
 - 2-methyl-4-(thien-2-ylethoxy)-6-[*N*-(5-carboxypyridin-2-yl)carbamoyl]benzofuran; and
 - 2-methyl-4-isobutoxy-6-[*N*-(thiazol-2-yl)carbamoyl]benzofuran;or a salt, solvate or pro-drug thereof.

7. (Original) A pharmaceutical composition comprising a compound according to any one of Claims 1 to 6, or a salt, pro-drug or solvate thereof, together with a pharmaceutically acceptable diluent or carrier.
8. (Currently Amended) A method of treating a disease mediated through glucokinase, comprising administering a compound according to any one of Claims 1 to 6 for use in the preparation of a medicament for treatment of a disease mediated through GLK.
9. (Currently Amended) A process for preparing a compound of formula (I); ~~as defined in Claim 1,~~



(I)

wherein:

Ring A is pyridin-2-yl or thiazol-2-yl; wherein said pyridin-2-yl or thiazol-2-yl is

optionally substituted on carbon by one or more groups selected from R⁴;

one of R¹ and R² is hydrogen and the other is hydrogen or C₁₋₄alkyl; wherein R¹ and R² are

optionally substituted on carbon by one or more groups selected from R⁵;

R³ is selected from C₁₋₄alkyl, C₁₋₄alkoxy, carbocyclyl, heterocyclyl, carbocyclyloxy and

heterocyclyloxy; wherein R³ is independently optionally substituted on carbon by

one or more groups selected from R⁶; and wherein if said heterocyclyl contains an

-NH- moiety that nitrogen is optionally substituted by C₁₋₄alkyl;

R⁴ is selected from halo, carboxy and C₁₋₄alkyl;

R⁵ and R⁶ are independently selected from halo, C₁₋₄alkyl, C₁₋₄alkoxy, N-(C₁₋₄alkyl)amino,

N,N-(C₁₋₄alkyl)₂amino, carbocyclyl, heterocyclyl, carbocyclyloxy, heterocyclyloxy

and carbocyclylidenyl; wherein R⁵ and R⁶ are independently optionally substituted


on carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH-

moiety that nitrogen is optionally substituted by C₁₋₄alkyl;

R⁷ is selected from halo, carboxy, methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, dimethylamino, diethylamino and N-methyl-N-ethylamino


or a salt, solvate or pro-drug thereof, which process (~~wherein variable groups are, unless otherwise specified, as defined in Claim 1~~) comprises:

Process 1): reacting an acid of formula (II):




(II)

or an activated derivative thereof; with a compound of formula (III); or



(III)

Process 2) for compounds of formula (I) wherein R⁴ is carboxy; deprotecting a compound of formula (III):




(III)

wherein R^xC(O)O- is an ester group;

and optionally thereafter if necessary or desirable:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- and/or
- ii) removing any protecting groups; and/or
- iii) forming a salt, solvate or pro-drug thereof, or a combination thereof.

10. A compound of formula (III); as defined in Claim 9



(III)

wherein:

R^xC(O)O- is an ester group;

Ring A is pyridin-2-yl or thiazol-2-yl; wherein said pyridin-2-yl or thiazol-2-yl is

optionally substituted on carbon by one or more groups selected from R⁴; and

one of R¹ and R² is hydrogen and the other is hydrogen or C₁₋₄alkyl; wherein R¹ and R² are optionally substituted on carbon by one or more groups selected from R⁵;

R³ is selected from C₁₋₄alkyl, C₁₋₄alkoxy, carbocyclyl, heterocyclyl, carbocyclyloxy and heterocyclyloxy; wherein R³ is independently optionally substituted on carbon by one or more groups selected from R⁶; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen is optionally substituted by C₁₋₄alkyl;

R⁴ is selected from halo, carboxy and C₁₋₄alkyl;

R⁵ and R⁶ are independently selected from halo, C₁₋₄alkyl, C₁₋₄alkoxy, N-(C₁₋₄alkyl)amino, N,N-(C₁₋₄alkyl)₂amino, carbocyclyl, heterocyclyl, carbocyclyloxy, heterocyclyloxy and carbocyclidenyl; wherein R⁵ and R⁶ are independently optionally substituted on carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen is optionally substituted by C₁₋₄alkyl; and

R⁷ is selected from halo, carboxy, methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, dimethylamino, diethylamino and N-methyl-N-ethylamino.